




Metal Ligand Bonding in Transition Metal Complexes

DATE

Valence Bond Theory (VBT):

Developed by Linus Pauling

According to VBT, the bonding in metal complexes arise when a filled ligand orbital containing a lone pair of electrons overlaps with a vacant hybrid orbital on the metal (ion) to give a coordinate covalent bond.



- VBT assumes the bonding between the metal (ion) and the ligands to be purely covalent.
- The hybrid orbitals that a metal (ion) uses in accepting the pair of electrons from the ligands are those that point in the directions of the ligands.

Salient features of VBT:

1. The central metal (ion) in the coordination compounds offers a number of vacant orbitals which is equal to the coordination number of the metal (ion) for the formation of coordinate bonds with the ligands.
2. The approximate atomic orbitals (s, p and d) of the metal hybridise to give a new set of orbitals of equivalent energy called the hybrid orbitals.
 - These hybrid orbitals are directed towards the ligands depending upon the geometry of the coordination complex.

3. The d-orbitals involved in the hybridization may be either inner $(n-1)d$ orbitals or outer nd orbitals. Thus, hybridisation in case of octahedral complex may involve d^2sp^3 or sp^3d^2 depending upon whether inner or outer d-orbitals are used for hybridization.


d^2sp^3 :: Inner orbital complex : Low spin complex
 sp^3d^2 : Outer orbital complex : High spin complex
 stability Order : $d^2sp^3 > sp^3d^2$

d-orbital involved for hybridization in octahedral complex = dx^2-y^2 and dz^2

for tetrahedral complex: sp^3 or sd^3 hybridization.




d-orbitals involved in sd^3 are dx^2-y^2 , dyz and dzx .

for square planar complexes: dsp^2 hybridization






4. The vacant hybrid orbitals of the metal (ion) overlap with the filled (or occupied) orbitals of the ligands and forms a coordinate bond.





Examples of complexes with C.N. 6 :




d^2sp^3
 Octahedral
 Inner orbital complex
 Diamagnetic



d^2sp^3
Octahedral
Inner orbital complex
Paramagnetic





d^2sp^3
Octahedral
Inner orbital complex
Diamagnetic




Inner orbital
Paramagnetic
classmate

d^2sp^3
octahedral


Complex Ion	O.S. of Metal	VSEC of Metal (ion)	Nature of Ligands	Valence Bond Electron Description	No. of Unpaired electrons
$[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$	+3	$3d^1$	Weak	<div style="text-align: center;"> $3d$ $4s$ $4p$ </div>  <p style="text-align: center;">d^2sp^3</p>	






- Octahedral
- Paramagnetic
- Outer orbital complex





- Octahedral
- Paramagnetic
- Outer orbital complex









- Octahedral, Paramagnetic
- Outer orbital complex






Tetrahedral
Paramagnetic



Tetrahedral
Diamagnetic



Square Planar
Diamagnetic





sp^3
Tetrahedral

Incorrect geometry

Correct Hybridization and Geometry:

Based ESR and X-ray structure determination studies: correct hybridization and geometry dsp^2 and square planar, respectively.



sp^2d hybridization

Square Planar

Limitations of VBT:

1. It fails to explain the colour and spectra of the complexes.
2. It fails to explain the differences between the measured and calculated magnetic moments, although the theory is successful in predicting the number of unpaired electrons in the complex.
3. It fails to explain why the magnetic moment values varies with temperature.
4. The theory fails to explain to the relative stabilities of the complexes.

5. It fails to explain why some complexes are labile while others are inert.

Labile Complexes: Those complexes in which one of the ligand can be easily displaced by another ligand.

Inert Complexes: Those complexes in which the displacement of ligands is slow.

6. It does not predict whether the 4-coordinate complex species will be square planar or tetrahedral.
e.g. $[\text{Cu}(\text{NH}_3)_4]^{2+}$

7. It considers $\text{Ni}(\text{acac})_2$ to be tetrahedral whereas in fact it has octahedral geometry.

8. The theory fails to explain why some metal complexes in a particular oxidation state are low spin (i.e. inner orbital complex) while some other complexes of the same metal (ion) in the same oxidation state are high spin (i.e. outer orbital complexes).

e.g. $[\text{CoF}_6]^{3-}$ High spin complex

$[\text{Co}(\text{NH}_3)_6]^{3+}$ Low spin complex